TDA-fall 2025


Extended Persistence (cont) + Reeb graphs Recap

· Paper chose: due October 17 (sorry for typo on web page!)
· Next assignment will be final

project proposal

6 Shorten office hours today


Extended Persistence Odd parts of persistence - points at Infinity -some Morse critical points (a_2, a_4)



Agarwol- Edelsbrumer-Haver-Wong 2006 3 Cohen-Steiner, Edels brunner, Herer 2009 Use relative homology to find other critical points, or get better paints. Relative homology Fix LCK Define Cp(K, L):= Cp(K)/Cp(L) 4 [x]={ 36Cp(K)] x+86Cp(L) Then maps extend to homology. Remember a month ago?


Let K = K 0 2x 0 260 2x 6 6 6 6 6 "Coned off

Theorem: $H_p(K,L) = H_p(K^*)$ for p>0 $A B_o(K,L) = B_o(K^*) - 1$




bits? ("cone off" $H_0(K, K^{a_1})$ $H_0(K,K^{a_8})$ $H_0(K, K^{a_6})$ $H_0(K, K^{a_5})$ $H_1(K,K^{a_8})$ $H_1(K,K^{a_6})$

Extended persistence Module regular persistence Hp (Kai) -> Hp (Ken)) $-- \longrightarrow H_p(K, K^{q_1})$ $H_p(K,K^{an}) \rightarrow$ relative Superlevel Set komology 3 kinds of points: o Ordinary: usual in PD o Relative: torn & Lie on way Journ · Extended: use both



Points in diagram; Hp(Kan) ---> Hp(Kan) -> Hp(K, Kan) ---> Hp(K, Kan) Ord: Starts & ends in 1st half, so (a,b) >> a < b Rel: Starts a ends in 2nd helf, (a,b) => 64a Ext: storts in 1st half => (a,b) enther on ch + (anan) -> ext Rel (ext)

On graphs



More generally; on manifolds, get symmetry Coulderta extended) · ordinary x relative

Under the hood: o Very Deautiful combination of Lefschetz of Poincare duality (that 2010 paper)

Some more algebraic connections Turner-Robins-Morgan





Recall: level sets

Gwen (PL) f: 1K/ >TR,


a level set is f'(a)

Ba: Consider f-1 ((-00,4]) A f-1 (4)






Prochce: What is f-1(10)



 $f^{-1}((-\infty,10))$

Answer just in case!



Sublevel set filton, 10]

Torus:


Kecall: A space is disconnected of I disjoint open Sets U, V with X=UOV L) otherwise, connected A convected component is a maximal subset that is anneated A contour 15 a connected component of a level set Q' How do contour changes?

Back to torangulation:

What shape, a how many components. (or recall for(10) a few slides back had 2 contours)

Equivalence relations of gusternts
A binery relation on a set T is an
equivalence relation if Ya, b, c e T · a ~ a [relexive] [symnetorc] 6 a ~ b ~ a if and about Thensitie] Equivalence dess of a under [a]= {xt] x~a} On a topological space with an equivalence relation, get quotient map q(a) = [a] (and U is open (=) q'(U) is open)

Reeb graph equivalence relation Given Function f: X->1R.



Define equiv. rel. by x ~ y H; · f(x)= f(y) = a · X + y are in some connected component of level set (16)

Reelo graphs · Let [x] donote the equivalence class of XEX. · The Keelo graph Rg = R(X,f) 1S the quotient space \times · Let D: X -> Kg. X Ho [x] be the quotient map

Viceness assumptions If input is "nice" (both X +f), then Ke is a graph. · Morse function on compact manifold oPL Rinchon of compact polyhedron 6 Offers ... Bad example: Non-Hausdorff Jisk/pt ()

topological signature Applications: Pascucci et falle in this figure: